

The minimum number of 4-cliques in a graph with triangle-free complement

Vladimir Nikiforov

Department of Mathematical Sciences
University of Memphis, Memphis, TN 38152

February 1, 2008

Abstract

Let $f(n, 4, 3)$ be minimum number of 4-cliques in a graph of order n with independence number 2. We show that

$$f(n, 4, 3) \leq \frac{1}{200}n^4 + O(n^3).$$

We also show if a graph of order n has independence number 2 and is close to regular then it has at least

$$\frac{1}{200}n^4 + o(n^4)$$

4-cliques.

1 Notation and conventions

Our notation and terminology are standard (see, e.g. [1]). In particular, all graphs are assumed to be defined on the vertex set $[n] = \{1, 2, \dots, n\}$. For any two adjacent vertices i and j we write $i \sim j$. We write $i \nsim j$ if i and j are distinct nonadjacent vertices. Given a vertex i , N_i denotes the set of its neighbors, d_i denotes its degree and t_i denotes the number of triangles containing it.

Given a graph G , $t_3(G)$ is the number of its triangles; $t_4(G)$ is the number of its 4-cliques; $t'_3(G)$ is the number of all induced subgraphs of order 3 and size 2; $t''_3(G)$ is the number of all induced subgraphs of order 3 and size 1, and $t'_4(G)$ is the number of all induced subgraphs of order 4 and size 5 (i.e. isomorphic to K_4 with one edge removed).

2 Introduction

Write $f(n, 4, 3)$ for the function

$$f(n, 4, 3) = \min\{t_4(G) : v(G) = n, \overline{G} \text{ is triangle-free}\}.$$

This function is a particular case of a more general function introduced by Erdős in [2]. In [3] we proved that

$$f(n, 4, 3) \leq \frac{1}{200}n^4 + o(n^4).$$

It happens that this estimate is tight under additional assumptions. We show that, in fact,

$$f(n, 4, 3) \leq \frac{1}{200}n^4 + O(n^3).$$

Moreover, let G be a graph of order n such that $\overline{G}$ is triangle-free. If G is almost regular, i.e., if

$$\sum_{i \in [n]} \left| d_i - \frac{2m}{n} \right| = o(n^2), \tag{1}$$

then we show that

$$t_4(G) \geq \frac{1}{200}n^4 + o(n^4).$$

3 Main results

Write $C_5[K_p]$ for the lexicographic product of the 5-cycle with the complete graph on p vertices. Recall that the vertex set of $G[K_p]$ is $v(G) \times v(K_p)$ and

$(i, x) \sim (j, y)$ iff $i \sim j$ or $x = y$. Observe that the complement of $C_5[K_p]$ is triangle-free and

$$t_4(C_5[K_p]) \geq 5 \left(\binom{2p}{4} - \binom{p}{4} \right) = \frac{25}{8}p^4 - \frac{35}{4}p^3 + \frac{55}{8}p^2 - \frac{5}{4}p.$$

Obviously, $f(n, 4, 3)$ is nondecreasing with respect to n . Let $5p$ be the smallest multiple of 5 which is not smaller than n . We have

$$\begin{aligned} f(n, 4, 3) &\leq 5 \left(\binom{2p}{4} - \binom{p}{4} \right) = \frac{25}{8}p^4 - \frac{35}{4}p^3 + \frac{55}{8}p^2 - \frac{5}{4}p \\ &\leq \frac{25}{8} \left(\frac{n+4}{5} \right)^4 - \frac{35}{4} \left(\frac{n+4}{5} \right)^3 + \frac{55}{8} \left(\frac{n+4}{5} \right)^2 - \frac{5}{4} \left(\frac{n+4}{5} \right) \\ &= \frac{1}{200}n^4 + \frac{1}{100}n^3 - \frac{17}{200}n^2 - \frac{13}{100}n + \frac{1}{5}. \end{aligned}$$

Our goal to the end of the paper is to prove the following assertion.

Theorem 1 *If a graph G of order n with independence number 2 satisfies (1) then*

$$t_4(G) \geq \frac{1}{200}n^4 + o(n^4).$$

The proof consists of the following steps:

- a) some lemmas establishing properties of graphs with triangle-free complement;
- b) deduction of a lower bound on $t_4(G)$ as a function of various graph parameters;
- c) reduction of the bound to a function of vertex degrees, $v(G)$ and $e(G)$;
- d) replacing the vertex degrees by their mean;
- e) minimizing the bound with respect to $e(G)$.

Lemma 1 *For any graph $G = G(n, m)$*

$$\sum_{i \sim j} d_i d_j \geq \frac{4m^3}{n^2}.$$

Proof Obviously, we can assume that G contains no isolated vertices. On the one hand, we have

$$2 \sum_{i \sim j} \frac{1}{\sqrt{d_i d_j}} \leq \sum_{i \sim j} \frac{1}{d_i} + \frac{1}{d_j} = \sum_{i \in [n]} \frac{d_i}{d_i} = n.$$

On the other hand, by the Cauchy-Schwarz inequality,

$$\left(\sum_{i \sim j} \sqrt{d_i d_j} \right) \left(\sum_{i \sim j} \frac{1}{\sqrt{d_i d_j}} \right) \geq m^2.$$

Therefore,

$$\sum_{i \sim j} \sqrt{d_i d_j} \geq \frac{2m^2}{n}.$$

Hence, by the Cauchy inequality,

$$\sum_{i \sim j} d_i d_j \geq \frac{1}{m} \left(\sum_{i \sim j} \sqrt{d_i d_j} \right)^2 \geq \frac{4m^3}{n^2}$$

completing the proof. $\square$

Let $G = G(n, m)$ be a graph with no independent set on 3 vertices (i.e., $\overline{G}$ is triangle-free). We shall prove a series of short lemmas which follow from this assumption.

Lemma 2

$$6t_3(G) = n^3 - 3n^2 + 2n + \sum_{i \in [n]} 3d_i^2 - 3d_i n + 3d_i.$$

Proof Indeed, this is an expanded form of a well known identity. Observe that

$$t'_3(G) + t''_3(G) = \frac{1}{2} \sum_{i \in [n]} d_i (n - 1 - d_i).$$

Hence,

$$\begin{aligned} 6t_3(G) &= 6 \binom{n}{3} - 6(t'_3(G) + t''_3(G)) \\ &= n^3 - 3n^2 + 2n - 3 \sum_{i \in [n]} d_i (n - 1 - d_i) \\ &= n^3 - 3n^2 + 2n + \sum_{i \in [n]} 3d_i^2 - 3nd_i + 3d_i. \end{aligned} \tag{2}$$

□

Lemma 3

$$2t_3(G) + t'_3(G) = (n-2)m - \binom{n}{3}. \quad (3)$$

Proof Indeed, trivially,

$$\binom{n}{3} = t_3(G) + t'_3(G) + t''_3(G).$$

On the other hand,

$$(n-2)m = 3t_3(G) + 2t'_3(G) + t''_3(G).$$

Subtracting the last two identities, we obtain (3). □

Lemma 4

$$8t_4(G) + 2t'_4(G) = \sum_{i \in [n]} (d_i - 2)t_i - \sum_{i \in [n]} \binom{d_i}{3}. \quad (4)$$

Proof Applying lemma 3 to $G[N_i]$ for any $i \in [n]$, and summing over all vertices, we obtain (4). □**Lemma 5**

$$t'_4(G) \leq \sum_{i \sim j} \binom{|N_i \cap N_j|}{2}.$$

Proof Obviously,

$$t'_4(G) = \sum_{i \sim j} e([N_i \cap N_j]) \leq \sum_{i \sim j} \binom{|N_i \cap N_j|}{2}.$$

□

Lemma 6 For any two nonadjacent vertices i and j

$$|N_i \cap N_j| = d_i + d_j - n + 2.$$

Proof For every k , if $k \sim j$ and $k \sim i$ then $\{i, j, k\}$ is a triangle in $\overline{G}$. Therefore,

$$n - 2 = |N_i \cup N_j| = d_i + d_j - |N_i \cap N_j|.$$

□

Lemma 7

$$\sum_{i \in [n]} d_i t_i = \sum_{i \sim j} d_i d_j - \frac{1}{2} \sum_{i \in [n]} d_i^2 - \frac{1}{2} \sum_{i \not\sim j} (d_i + d_j) |N_i \cap N_j|.$$

Proof Let i be any vertex. Obviously,

$$\sum_{j \sim i} d_j = d_i + 2t_i + \sum_{j \not\sim i} |N_i \cap N_j|$$

hence, multiplying both sides by d_i , we get

$$d_i \sum_{i \sim j} d_j = d_i^2 + 2d_i t_i + d_i \sum_{j \not\sim i} |N_i \cap N_j|$$

and summing over all vertices, we obtain

$$2 \sum_{i \sim j} d_i d_j = \sum_{i \in [n]} d_i^2 + 2 \sum_{i \in [n]} d_i t_i + \sum_{j \not\sim i} (d_i + d_j) |N_i \cap N_j|.$$

□

Proof of Theorem 1 By lemma 4 and lemma 5,

$$\begin{aligned} 8t_4(G) &= -2t'_4(G) + \sum_{i \in [n]} (d_i - 2) t_i - \sum_{i \in [n]} \binom{d_i}{3} \\ &\geq -2 \sum_{i \not\sim j} \binom{|N_i \cap N_j|}{2} - 6t_3(G) + \sum_{i \in [n]} d_i t_i - \sum_{i \in [n]} \binom{d_i}{3}. \end{aligned}$$

Hence, by lemma 6 and lemma 7,

$$\begin{aligned}
8t_4(G) &\geq - \sum_{i \sim j} (d_i + d_j - n + 2)(d_i + d_j - n + 1) - 6t_3(G) - \sum_{i \in [n]} \binom{d_i}{3} \\
&\quad + \sum_{i \sim j} d_i d_j - \frac{1}{2} \sum_{i \in [n]} d_i^2 - \frac{1}{2} \sum_{i \sim j} (d_i + d_j)(d_i + d_j - n + 2) \\
&= \sum_{i \sim j} d_i d_j - \frac{1}{2} \sum_{i \sim j} (d_i + d_j - n + 2)(3d_i + 3d_j - 2n + 2) \\
&\quad - \frac{1}{6} \sum_{i \in [n]} (d_i^3 - 3d_i^2 + 2d_i) - \frac{1}{2} \sum_{i \in [n]} d_i^2 - 6t_3(G).
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
8t_4(G) &\geq \sum_{i \sim j} d_i d_j - \frac{1}{2} \sum_{i \sim j} (d_i + d_j - n + 2)(3d_i + 3d_j - 2n + 2) \quad (5) \\
&\quad - \frac{1}{6} \sum_{i \in [n]} (d_i^3 + 2d_i) - 6t_3(G).
\end{aligned}$$

We also find that

$$\begin{aligned}
&\sum_{i \sim j} (d_i + d_j - n + 2)(3d_i + 3d_j - 2n + 2) \\
&= \sum_{i \sim j} (3d_i^2 + 6d_i d_j - 5nd_i + 8d_i + 3d_j^2 - 5nd_j + 8d_j + 2n^2 - 6n + 4) \\
&= \sum_{i \sim j} 6d_i d_j + \sum_i (3d_i^2(n-1-d_i) - 5nd_i(n-1-d_i) + 8d_i(n-1-d_i)) + \\
&\quad + \sum_{i \in [n]} (n^2 - 3n + 2)(n-1-d_i) \\
&= \sum_{i \sim j} 6d_i d_j + \sum_{i \in [n]} (8nd_i^2 - 3d_i^3 - 11d_i^2 - 6n^2 d_i + 16nd_i - 10d_i) \\
&\quad + n^4 - 4n^3 + 5n^2 - 2n.
\end{aligned}$$

Applying this equality to (5) and afterwards bounding $6t_3$ by lemma 2, we

get

$$\begin{aligned}
8t_4(G) &\geq \sum_{i \sim j} d_i d_j - 3 \sum_{i \not\sim j} d_i d_j - \frac{1}{2} (n^4 - 4n^3 + 5n^2 - 2n) - \frac{1}{6} \sum_i (d_i^3 + 2d_i) \\
&\quad - 6t_3 + \sum_i \left(\frac{3}{2} d_i^3 - 4n d_i^2 + 3n^2 d_i + \frac{11}{2} d_i^2 - 8n d_i + 5d_i \right) \\
&= \sum_{i \sim j} d_i d_j - 3 \sum_{i \not\sim j} d_i d_j - \frac{1}{2} n^4 + n^3 + \frac{1}{2} n^2 - n + \\
&\quad + \sum_i \left(\frac{4}{3} d_i^3 - 4n d_i^2 + 3n^2 d_i + \frac{5}{2} d_i^2 - 5n d_i + \frac{5}{3} d_i \right)
\end{aligned}$$

On the other hand, note that

$$\begin{aligned}
\sum_{i \sim j} d_i d_j - 3 \sum_{i \not\sim j} d_i d_j &= 4 \sum_{i \sim j} d_i d_j - \frac{3}{2} \left(\sum_{i \in [n]} d_i \right)^2 + \frac{3}{2} \sum_{i \in [n]} d_i^2 \\
&= 4 \sum_{i \sim j} d_i d_j - 6m^2 + \frac{3}{2} \sum_{i \in [n]} d_i^2
\end{aligned}$$

Thus,

$$\begin{aligned}
8t_4(G) &\geq 4 \sum_{i \sim j} d_i d_j - 6m^2 - \frac{1}{2} n^4 + n^3 + \frac{1}{2} n^2 - n + \\
&\quad + \sum_{i \in [n]} \left(\frac{4}{3} d_i^3 - 4n d_i^2 + 3n^2 d_i + 4d_i^2 - 5n d_i + \frac{5}{3} d_i \right)
\end{aligned}$$

Dropping the low order terms, we see that

$$8t_4 \geq 4 \sum_{i \sim j} d_i d_j - 6m^2 - \frac{1}{2} n^4 + \sum_{i \in [n]} \left(\frac{4}{3} d_i^3 - 4n d_i^2 + 3n^2 d_i \right) + O(n^3).$$

Due to (1), we find that

$$\begin{aligned}
8t_4 &\geq 4m \frac{4m^2}{n^2} - 6m^2 - \frac{1}{2} n^4 + \frac{4}{3} \frac{8m^3}{n^2} - 4n \frac{4m^2}{n} + 6n^2 m + O(n^3) \\
&= \frac{80}{3} \frac{m^3}{n^2} - 22m^2 + 6mn^2 - \frac{1}{2} n^4 + O(n^3).
\end{aligned}$$

Since the expression

$$\frac{80}{3} \frac{m^3}{n^2} - 22m^2 + 6mn^2$$

attains its minimum at

$$m = \frac{3}{10}n^2 + o(n^2),$$

the desired inequality follows. $\square$

Acknowledgment. The author is grateful to Andrew Thomason for his encouraging attention.

References

- [1] B. Bollobás, *Modern Graph Theory*, Graduate Texts in Mathematics, **184**, Springer-Verlag, New York (1998), xiv+394 pp.
- [2] P. Erdős, On the number of complete subgraphs contained in certain graphs, *Publ. Math. Inst. Hungar. Acad. Sci.* 7(1962) 459-464.
- [3] V. Nikiforov, On the minimal number of k -cliques in graphs with restricted number of independence, *Comb. Prob. Comp.* **10** (2001), 361-366.